



Mycoplasma alkalescens demonstrated in bronchoalveolar lavage of cattle in Denmark

Kokotovic, Branko; Friis, Niels F.; Ahrens, Peter

Published in:
Acta Veterinaria Scandinavica (Online Edition)

Link to article, DOI:
[10.1186/1751-0147-49-2](https://doi.org/10.1186/1751-0147-49-2)

Publication date:
2007

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Kokotovic, B., Friis, N. F., & Ahrens, P. (2007). Mycoplasma alkalescens demonstrated in bronchoalveolar lavage of cattle in Denmark. *Acta Veterinaria Scandinavica (Online Edition)*, 49, 2. <https://doi.org/10.1186/1751-0147-49-2>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Brief communication

Open Access

***Mycoplasma alkalescens* demonstrated in bronchoalveolar lavage of cattle in Denmark**

Branko Kokotovic*, Niels F Friis and Peter Ahrens

Address: National Veterinary Institute, New Technical University of Denmark, Bülowsvej 27, DK-1790 Copenhagen V, Denmark

Email: Branko Kokotovic* - bko@vet.dtu.dk; Niels F Friis - nff@vet.dtu.dk; Peter Ahrens - pa@vet.dtu.dk

* Corresponding author

Published: 04 January 2007

Received: 26 September 2006

Acta Veterinaria Scandinavica 2007, **49**:2 doi:10.1186/1751-0147-49-2

Accepted: 04 January 2007

This article is available from: <http://www.actavetscand.com/content/49/1/2>

© 2007 Kokotovic et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Mycoplasma alkalescens is an arginine-metabolizing mycoplasma, which has been found in association with mastitis and arthritis in cattle. Routine bacteriological examination of 17 bronchoalveolar lavage samples from calves with pneumonia in a single herd in Denmark, identified *M. alkalescens* in eight samples. The organism was found as a sole bacteriological findings in five of the samples as well as in combination with *Mannheimia haemolytica*, *Haemophilus somni* and *Salmonella* Dublin. This is the first report of isolation of *M. alkalescens* in Denmark.

Findings

Mycoplasma alkalescens is a bovine mycoplasma species, which was originally isolated from the nasal cavity of cattle in Australia [1]. Like many other mycoplasmas, *M. alkalescens* is a normal inhabitant of the upper respiratory tract, but it has also been associated with disease. *M. alkalescens* has mostly been implicated in mastitis in cattle. It has been isolated from bulk tank milk samples, as well as from outbreaks and sporadic cases of clinical mastitis [2-5]. Furthermore, *M. alkalescens* has been isolated from cases of severe arthritis, and its ability to induce joint lesions has been confirmed under experimental conditions [6,7]. Rosenfeld & Hill [8] isolated *M. alkalescens* in pure culture from abomasum and lung of an aborted bovine foetus, while Lamm et al. [9] found *M. alkalescens* in association with otitis in calves. Finally, *M. alkalescens* has occasionally been found in association with disorders of the respiratory tract [10,11].

Recently, 17 bronchoalveolar lavage samples from calves suffering from pneumonia in a single herd in Denmark were submitted, on two occasions, to the National Veteri-

nary Institute for laboratory examinations. The samples were examined for the presence of bacterial pathogens, bovine respiratory mycoplasmas, as well as for the presence of bovine respiratory syncytial virus (BRSV), bovine coronavirus and parainfluenza-3 virus (PI-3). The samples were not examined for infectious bovine rhinotracheitis virus, as Denmark is considered free of this infection. Thereby, eight arginine-metabolizing mycoplasmas were isolated. In the present study the identification of the isolates as *M. alkalescens* is presented. This is the first report of isolation of this species in Denmark.

Bacteriological examination was performed according to standard laboratory procedures. Examination for BRSV, bovine coronavirus and PI-3 was performed using an indirect sandwich-ELISA assay [12]. Isolation of mycoplasmas was performed according to standard laboratory procedures using a Hayflick's type of medium enriched with arginine [13], with and without addition of 5% of rabbit hyperimmune antiserum against *Mycoplasma bovirhinis*. The isolates were filtered through 0.45 µm membrane filters (Millipore), cloned and submitted to serologic identi-

fication, which was performed by the disc growth inhibition test (DGI) with 6 mm filter paper discs prepared with rabbit hyperimmune antisera against *M. alkalescens* PG51^T (ATCC 29103; NCTC 10135), *Mycoplasma arginini* G230^T (ATCC 23838; NCTC 10129) and *Mycoplasma canadense* 275C^T (ATCC 29418; NCTC 10152), as well as by the indirect epi-immunofluorescence test (IF) on colonies on solid medium. Molecular identification of the isolates was performed by amplification and sequencing of their 16S rRNA genes by using universal primers [14].

Laboratory examinations revealed the presence of several bacterial pathogens in 15 of the 17 bronchoalveolar lavage samples, while no pathogenic bacteria were detected in the two remaining samples. *Pasteurella multocida*, *Mannheimia haemolytica*, *Histophilus somni* and *Salmonella* Dublin were found in three, three, one and one samples, respectively. *Mycoplasma bovirhinis* was found in five samples, while arginin-metabolizing mycoplasmas were found in eight samples. Two *P. multocida*, three *M. bovirhinis* and five arginin-metabolizing mycoplasma isolates were found as sole bacteriological findings in 10 respective samples. From five samples multiple bacterial species were isolated: *M. bovirhinis* was found once in combination with *P. multocida* and once in combination with *M. haemolytica*, while arginin-metabolizing mycoplasmas were found once in combination with *M. haemolytica*, once in combination with *S. Dublin* and once in combination with *M. haemolytica* and *H. somni*. No viral pathogens were detected in any of the analysed samples.

Further examinations on arginin-metabolizing mycoplasmas showed that all eight isolates could be serologically identified as *M. alkalescens* and that they were clearly dif-

ferent from the two other arginine-degrading species that are commonly found in cattle, *M. canadense* and *M. arginini* (Table 1). The reactions were clear-cut for the DGI tests showing approximately 3–5 mm broad zones of total or nearly total inhibition around the disc. In the IF test some minor cross reactions were noted when three of the isolates and the type strain of *M. alkalescens* PG51^T were tested with anti-*M. canadense* 275C^T hyperimmune serum. The 16S rDNA sequences of the analysed isolates were identical to each other and to the 16S rDNA sequence of the type strain *M. alkalescens* PG51^T (GenBank accession no. [U44764](#); [15]), which corroborated the serological identification of the isolates as *M. alkalescens*.

Previous studies have shown that *H. somni*, *M. haemolytica* and *P. multocida* are the bacteria that are most commonly associated with bronchopneumonia in calves in Denmark, although they are also part of the normal bacterial flora of the respiratory tract [16,17]. Also, *S. Dublin*, which is predominantly found in cattle, is capable of causing pneumonia in calves. Isolation of these bacterial species in the submitted samples most likely indicate their role as the primary cause of respiratory disease in the herd, probably in combination with unidentified environmental factors. In addition to the bacteria, *M. bovirhinis* and *M. alkalescens* were also isolated. *M. bovirhinis* is commonly found as a part of the normal respiratory tract microflora of cattle and is not considered to be pathogenic. *M. alkalescens*, however, has been found in association with disorders of the respiratory tract but its role as a respiratory pathogen remains equivocal. Experimental infections have demonstrated that *M. alkalescens* has the ability to colonize lung tissue [18], but the two strains used in the study apparently failed to produce pneumonia [18]. In the present study, we found *M. alkalescens* in pure culture

Table 1: Serological identification of eight mycoplasma field isolates from cattle by DGI and IF test using type strain antisera against *Mycoplasma alkalescens*, *Mycoplasma arginini* and *Mycoplasma canadense*.

Mycoplasma strains	Antiserum					
	<i>M. alkalescens</i> PG51 ^T		<i>M. arginini</i> G230 ^T		<i>M. canadense</i> 275C ^T	
	DGI ^a	IF ^b	DGI ^a	IF ^b	DGI ^a	IF ^b
<i>M. alkalescens</i> PG51 ^T	5	+	0	-	0	?
Field strain MK19/02	≥ 3	+	0	-	0	-
Field strain MK20/02	≥ 3	+	0	-	0	?
Field strain MK21/02	≥ 3	+	0	-	0	-
Field strain MK49/02*	≥ 3	+	0	-	0	?
Field strain MK51/02	≥ 3	+	0	-	0	-
Field strain MK52/02*	≥ 3	+	0	-	0	-
Field strain MK53/02	≥ 3	+	0	-	0	-
Field strain MK56/02*	≥ 3	+	0	-	0	?
<i>M. arginini</i> G230 ^T	0	-	5	+	0	-
<i>M. canadense</i> 275C ^T	0	-	0	-	5	+

^a zone of inhibition in mm; ^b distinct FITC colour of stained colonies; ? inconclusive results; * examined by 16S rDNA sequencing.

in five of the analysed bronchoalveolar lavage samples. This finding is, however, not sufficient to warrant a role of *M. alkalescens* as a cause of bronchopneumonia, since we also found two samples containing only a non-pathogenic *M. bovirhinis*, and two samples without any bacterial or viral pathogens, despite the fact that they derived from calves with clinical signs of a respiratory disease. The failure to detect respiratory tract pathogens in these samples may be due to i.e., their absence in a particular disease stadium when sampling took a place or due to antibiotic treatment. Taking into the consideration the overall bacteriological findings of this study, it seems likely that *M. alkalescens* may have had a role either as a secondary invader or as an opportunistic pathogen rather than suggesting a causal role of the organism in the pneumonia complex.

M. alkalescens is regarded as one of the most common causative agents of mastitis [19]. So, with demonstration of the organism in a Danish cattle herd, a further member of *Mycoplasma* is added to the group of bovine mastitis-inducing microorganisms in Denmark. The only mycoplasma species isolated from clinical outbreaks of mastitis in Denmark so far has been *Mycoplasma bovis* [20], while other mastitis-inducing species, *Mycoplasma bovigenitalium* and *M. canadense*, have been isolated only from the respiratory and the genital tract and semen samples [21]. Further investigations are needed in order to determine the prevalence of *M. alkalescens* in the Danish cattle population and, indeed, to draw a firm conclusions on its importance in disease conditions other than mastitis.

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

BK carried out bacteriological examination of samples, isolation and cloning of mycoplasmas and drafted the manuscript.

NFF carried out identification of mycoplasma isolates by DGI test and epi-immunofluorescence.

PA carried out molecular identification of mycoplasma isolates.

All authors read and approved the final manuscript.

Acknowledgements

Authors thank Ulla Amtoft for skilful technical assistance.

References

- Hudson JR, Etheridge JR: **A new type of pleuropneumonia-like microorganism (PPLO) from the nose of cattle.** *Aust Vet J* 1963, **39**:1-5.

- Brookbanks EO, Carter ME, Holland JTS: **Mycoplasma mastitis.** *NZ Vet J* 1969, **17**:179-180.
- Dellinger JD, Jasper DE, Ilic M: **Characterization studies on mycoplasmas isolated from bovine mastitis and the bovine respiratory tract.** *Cornell Vet* 1977, **67**:351-360.
- Jasper DE: **Prevalence of mycoplasmal mastitis in the Western States.** *California Vet* 1980, **34**:24-26.
- Jasper DE: **The role of Mycoplasma in bovine mastitis.** *J Amer Vet Med Assoc* 1982, **181**:158-162.
- Bennett RH, Jasper DE: **Mycoplasma alkalescens-induced arthritis in dairy calves.** *J Amer Vet Med Assoc* 1978, **172**:484-488.
- Whithear KG: **Isolation of Mycoplasma alkalescens from cases of polyarthritis in embryo-transplant calves.** *Aust Vet J* 1983, **60**:191-192.
- Rosenfeld LE, Hill MWM: **The isolation of Mycoplasma alkalescens from an aborted bovine foetus.** *Aust Vet J* 1980, **56**:350.
- Lamm CG, Munson L, Thurmond MC, Barr BC, George LW: **Mycoplasma otitis in California calves.** *J Vet Diagn Invest* 2004, **16**:397-402.
- Hewicker-Trautwein M, Feldmann M, Kehler W, Schmidt R, Thiede S, Seeliger F, Wohlsein P, Ball HJ, Buchenau I, Spersger J, Rosengarten R: **Outbreak of pneumonia and arthritis in beef calves associated with Mycoplasma bovis and Mycoplasma californicum.** *Vet Rec* 2002, **151**:699-703.
- Ayling RD, Bashiruddin SE, Nicholas RAJ: **Mycoplasma species and related organisms isolated from ruminants in Britain between 1990 and 2000.** *Vet Rec* 2004, **155**:413-416.
- Uttenthal A, Jensen NPB, Blom JY: **Viral aetiology of enzootic pneumonia in Danish dairy herds: Diagnostic tools and epidemiology.** *Vet Rec* 1996, **139**:114-117.
- Kobisch M, Friis NF: **Swine mycoplasmosis.** *Rev Sci Tech Off Int Epiz* 1996, **15**:1569-1605.
- Weisburg WG, Barns SM, Pelletier DA, Lane DJ: **16S ribosomal DNA amplification for phylogenetic study.** *J Bacteriol* 1991, **173**:697-703.
- Pettersson B, Uhlen M, Johansson K-E: **Phylogeny of some mycoplasmas from ruminants based on 16S rRNA sequences and definition of a new cluster within the hominis group.** *Int J Syst Bacteriol* 1996, **46**:1093-1098.
- Tegtmeier C, Uttenthal A, Friis NF, Jensen NE, Jensen HE: **Pathological and microbiological studies on pneumonic lungs from Danish calves.** *J Vet Med* 1999, **46**:693-700.
- Watts JL, Yancey RJ, Sarah JR, Salmon SA, Case CA: **A 4-year survey of antimicrobial susceptibility trends for isolates from cattle with bovine respiratory disease in North America.** *J Clin Microbiol* 1994, **32**:725-731.
- Gourlay RN, Howard CJ, Thomas LH, Wyld SG: **Pathogenicity of some Mycoplasma and Acholeplasma species in the lungs of gnotobiotic calves.** *Res Vet Sci* 1979, **27**:233-237.
- Gonzalez RN, Wilson DJ: **Mycoplasma mastitis in dairy herds.** *Vet Clin North Am Food Anim Pract* 2003, **19**:199-221.
- Friis NF: **Mycoplasma bovis-induced mastitis in cattle in Denmark.** *Nord Vet Med* 1984, **36**:324-325.
- Friis NF, Blom E: **Isolation of Mycoplasma canadense from bull semen.** *Acta Vet Scand* 1983, **24**:315-317.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

